



Comparing Muscle Oxygenation and Peak Blood Lactate Concentration in Division 1 College Middle-Distance Athletes: A Speed Reserve Ratio Analysis

Braeden Kraft¹, Ayla Kraft², Evan E. Schick¹, Kurt A. Escobar¹, Kevin A. Valenzuela^{1,*}  and Joshua A. Cotter¹

¹Department of Kinesiology, California State University Long Beach, Long Beach, CA, USA

²Department of Biology, California State University Long Beach, Long Beach, CA, USA

Abstract:

Introduction: Middle-distance running requires a balance of aerobic and anaerobic energy systems, necessitating efficient physiological profiling for optimized training. Performance biomarkers, such as the Speed Reserve Ratio (SRR), have been used to improve training and performance but may not always be practical for routine assessment. Other biomarkers, such as skeletal muscle oxygenation (SmO_2) and blood lactate concentration ([La]bmax), may be more cost-effective and easier to integrate into training programs than the SRR. This study investigates the relationship between SmO_2 and [La]bmax with SRR in NCAA Division I middle-distance runners to determine whether these physiological markers could serve as viable alternatives for athlete profiling.

Methods: Male NCAA Division I athletes ($n = 11$) completed a series of sprint trials and incremental treadmill tests to assess SmO_2 and blood lactate responses.

Results: The change in skeletal muscle oxygenation (ΔSmO_2) exhibited a significant positive correlation with SRR ($r = 0.656$, $p = 0.028$), while Reoxygenation Hemoglobin Slope (ROHS) displayed a negative correlation ($r = -0.644$, $p = 0.033$), suggesting that oxygen utilization and recovery kinetics differentiate athletes with higher anaerobic or aerobic capacities.

Discussion: These findings highlight the potential of SmO_2 -derived metrics as accessible, noninvasive alternatives for profiling aerobic-anaerobic balance in middle-distance runners, particularly in situations where SRR measurement is impractical or unavailable. Future research should explore how these markers integrate into training models, their predictive validity across different competition levels, and their application in optimizing individualized athlete development programs.

Conclusion: Integrating noninvasive SmO_2 monitoring can be a beneficial addition to training programs for enhancing performance outcomes in middle-distance runners.

Keywords: Middle distance running, Physiological profiling, NCAA Division I, Speed reserve ratio, Blood lactate concentration.

© 2026 The Author(s). Published by Bentham Open.

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: <https://creativecommons.org/licenses/by/4.0/legalcode>. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



CrossMark

Received: July 23, 2025

Revised: September 12, 2025

Accepted: October 15, 2025

Published: January 26, 2026



Send Orders for Reprints to
reprints@benthamscience.net

*Address correspondence to this author at the Department of Kinesiology, California State University Long Beach, Long Beach, CA, USA; E-mail: kevin.valenzuela@csulb.edu

Cite as: Kraft B, Kraft A, Schick E, Escobar K, Valenzuela K, Cotter J. Comparing Muscle Oxygenation and Peak Blood Lactate Concentration in Division 1 College Middle-Distance Athletes: A Speed Reserve Ratio Analysis. Open Sports Sci J, 2026; 19: e1875399X431328251130185212. <http://dx.doi.org/10.2174/011875399X431328251130185212>

1. INTRODUCTION

Middle-distance running is a unique discipline in track and field that relies on a close balance of energy contributions from both aerobic and anaerobic energy systems [1, 2]. This dual energetic demand helps explain why these events are considered the “middle ground” of aerobic and anaerobic abilities, where adaptations to both energy systems can influence performance. These varying energy demands lead to a physiologically diverse group of competitors, with some athletes achieving success through higher aerobic capacity and others through anaerobic power [3, 4]. This diversity means that a one-size-fits-all training approach may not be effective. Profiling helps illuminate an athlete’s strengths and, more importantly, their weaknesses, allowing coaches to individualize training that maintains an athlete’s strengths and improves the areas where they are lacking. This is especially relevant in events like the 800 meters, where some of the world’s best athletes are successful due to higher maximal sprint speed and anaerobic speed reserve, enabling them to respond to late-race surges and pace changes [3]. Identifying these traits can help coaches align training with the specific demands of the event. In addition, profiling has been used to guide targeted training interventions—athletes who excel aerobically but underperform anaerobically can improve performance by training their anaerobic system more directly [5]. Researchers and coaches can categorize an athlete’s abilities using different physiological profiling techniques. A physiological profile refers to the comprehensive characterization of an athlete’s aerobic, anaerobic, and neuromuscular/mechanical capabilities, reflecting their unique bioenergetic and biomechanical strengths and limitations that influence performance [6-8].

Middle-distance runners tend to vary in bioenergetic ability, highlighting the need for accurate profiling of biomarkers that can improve understanding of an athlete’s unique strengths and weaknesses [9]. Traditional profiling metrics, like $\text{VO}_{2\text{max}}$ and lactate threshold, have been shown to be insufficient in predicting middle-distance race performance and fail to encompass an athlete’s anaerobic abilities [3, 4]. A more recent metric used in middle-distance profiling is the Speed Reserve Ratio (SRR), which addresses the limitations of other metrics by measuring both an athlete’s aerobic and anaerobic capacity. The SRR is calculated as the ratio of an athlete’s maximum sprint speed (MSS) to their maximum aerobic speed (MAS). Higher SRR values indicate greater anaerobic power, while lower SRR values indicate stronger aerobic capacity [10]. By capturing both speed domains of performance, SRR yields a more comprehensive bioenergetic profile [11].

Despite its utility as a profiling metric, obtaining accurate SRR values can be difficult due to the specialized equipment required for testing, such as high-speed treadmills, metabolic carts, and laser timing systems. Practical limitations of accurately measuring SRR have prompted interest in other noninvasive biomarkers that could serve as alternatives. Consequently, newer

techniques are being explored because of their portability, relatively low cost, sensitivity to training adaptations, and demonstrated links to middle-distance performance [12-20].

Muscle oxygen saturation (SmO_2), which refers to the balance between oxygen delivery and utilization within skeletal muscle and is commonly assessed noninvasively using near-infrared spectroscopy (NIRS), is one alternative metric used to profile athletes. Several SmO_2 -derived metrics have correlated with athletic performance outcomes in both sprint and endurance athletes. These metrics include SmO_2 delta (ΔSmO_2), muscle deoxygenation slope (MdO_2 slope), reoxygenation slope (ROHS), and the inflection point (IP). ΔSmO_2 is calculated as the percent change from baseline to minimum SmO_2 during exercise, reflecting local oxygen extraction and oxidative efficiency [16]. MdO_2 slope represents the initial rate of desaturation during the onset of exercise, indicating the speed of aerobic engagement [17]. IP is the breakpoint in the SmO_2 curve during exercise, corresponding to systemic lactate threshold and the shift toward anaerobic metabolism [18]. ROHS is the rate of SmO_2 recovery post-exercise, assessing the muscle’s capacity for oxygen restoration [19]. Although these SmO_2 metrics have proven useful in athletics, their ability to profile and distinguish physiological subgroups in middle-distance runners remains unexplored.

Peak post-exercise blood lactate concentration ($[\text{La}]_{\text{bmax}}$), defined as the highest blood lactate value measured after exercise, has also been associated with both middle-distance performance and an athlete’s anaerobic capacity [20, 21]. Athletes with higher $[\text{La}]_{\text{bmax}}$ values tend to demonstrate greater anaerobic power and sprint performance, while lower values may indicate improved lactate clearance and more efficient aerobic metabolism [14]. This balance between lactate production and clearance is relevant in middle-distance performance, where both anaerobic and aerobic systems are significantly taxed. However, despite the utility of $[\text{La}]_{\text{bmax}}$ in identifying general metabolic capabilities, its interpretability is constrained by physiological and methodological factors [22, 23]. In addition, the correlational strength between $[\text{La}]_{\text{bmax}}$ and SRR has yet to be examined.

Given these considerations, this study aims to determine whether SmO_2 -derived metrics and/or $[\text{La}]_{\text{bmax}}$ can serve as practical, noninvasive alternatives to SRR for profiling the aerobic-anaerobic balance in NCAA Division I middle-distance runners. We hypothesized that SmO_2 -derived metrics, including the percent change in muscle oxygenation (ΔSmO_2), the muscle deoxygenation slope (MdO_2 slope), the inflection point (IP), and the reoxygenation slope (ROHS), would significantly correlate with SRR, offering insight into an athlete’s balance of aerobic and anaerobic capabilities. Specifically, we expected that higher ΔSmO_2 values and steeper MdO_2 slopes would be associated with lower SRR values, reflecting greater aerobic efficiency. We also hypothesized

that a higher IP would occur in athletes with lower SRR values, indicating the ability to sustain higher workloads before shifting toward anaerobic metabolism. Additionally, we predicted that a steeper ROHS would correlate with lower SRR values, signifying more efficient oxygen recovery following maximal exercise. Finally, we expected that elevated [La]bmax would correlate with higher SRR values, reinforcing its role as an indicator of anaerobic power.

2. METHODOLOGY

2.1. Study Design

This cross-sectional study examined relationships among SmO_2 metrics, [La]bmax, and SRR in NCAA Division I middle-distance runners. All testing took place at the Movement Science Lab and Jack Rose Track at CSULB. To minimize fatigue and ensure data quality, testing was divided into three separate days, with a minimum of 48 hours of recovery provided between sessions. Day 1 included baseline anthropometrics and a familiarization trial. Day 2 assessed MSS via sprint tests on the outdoor track. Day 3 consisted of an incremental treadmill protocol to determine MAS and record SmO_2 via NIRS, with post-test blood samples collected for [La]bmax. Distributing assessments in this manner reduced carry-over fatigue and ensured maximal effort in both anaerobic and aerobic tests. Ethical clearance was granted by the Institutional Review Board (IRB) of CSU Long Beach, ensuring adherence to ethical standards in research involving human subjects.

2.2. Procedures

All testing procedures were performed under controlled conditions, with standardized warm-ups and cooldowns implemented to maintain participant consistency.

2.2.1. Day 1: Anthropometric Assessments and Familiarization Trial

The first session focused on baseline measurements and participant familiarization with the experimental protocol. Upon arrival, height and body mass were recorded using a calibrated Health O Meter 500KL stadiometer and digital scale. Body composition was assessed via seven-site skinfolds (triceps, subscapular, chest, midaxillary, suprailiac, abdomen, thigh), with measurements taken using a Lange skinfold caliper and calculated using the Jackson and Pollock equation [24]. A single investigator performed all body composition assessments to minimize measurement variability. For the familiarization test, participants completed a self-paced three-mile treadmill run while wearing a reusable face mask (COSMED, Rome, Italy). The Train.RED FYER NIRS device (Train.Red, Elst, Netherlands) was affixed to the vastus lateralis muscle to assess SmO_2 . This familiarization session aimed to minimize learning effects in subsequent trials and ensure participants could perform at maximal effort without interference from unfamiliar equipment.

Due to equipment constraints, six of the 11 participants completed their trials on a Cybex 751T treadmill, and the remaining five used a Woodway ELG treadmill. Although both models are widely used in exercise testing, minor differences in belt speed calibration, shock absorption, and surface characteristics may introduce variability.

2.2.2. Day 2: Maximum Sprint Speed Testing

The second session focused on testing for MSS on the Jack Rose synthetic track at California State University, Long Beach. Participants completed a standardized warm-up routine, which included five minutes of low-intensity jogging, dynamic stretching, and strides. Each participant then performed three 50-meter sprints on a flat synthetic track, with at least three minutes of passive recovery between attempts to mitigate fatigue. A 50 m all-out sprint was used to capture maximal sprint speed, as this distance reliably elicits near-maximum velocity in athletic populations [25]. Sprint velocity was measured using a Brower TCI Timing System, with laser timing gates positioned at 40 meters and 50 meters to capture peak speed. The highest recorded velocity across the three trials was designated as the participant's MSS.

2.2.3. Day 3: Max Aerobic Speed Testing

The final session consisted of an incremental treadmill test designed to assess MAS, SmO_2 , and [La]bmax. Before testing, participants were fitted with a Polar H10 heart rate monitor, a COSMED face mask connected to a metabolic analyzer (COSMED, Rome, Italy), and the NIRS optode secured to the vastus lateralis muscle with adhesive patches and an elastic bandage. Baseline SmO_2 values were then recorded while participants remained seated for five minutes, providing a resting reference point [25].

Participants then completed a standardized ten-minute warm-up, beginning with five minutes of running at 14 km/h, followed by dynamic stretching. The treadmill test protocol commenced at 14 km/h, increasing by 1 km/h every minute until 19 km/h. Beyond this point, increments decreased to 0.5 km/h per minute until 21 km/h, and then to 0.2 km/h per minute until voluntary exhaustion [12]. SmO_2 was continuously monitored to assess oxygen extraction and utilization in working muscle tissue.

Post-exercise blood lactate concentration in millimoles per liter (mmol/L) was measured to assess anaerobic energy system engagement. Blood samples were collected from the earlobe at one, three, five, and seven minutes following the treadmill test using a Lactate Scout 2 analyzer (EKF Diagnostics) [12, 26]. The highest value recorded during this period was designated as [La]bmax, representing peak lactate accumulation in response to exercise. The timing of lactate sampling was intended to ensure that peak concentrations were captured within the expected post-exercise window [26]. Although post-exercise blood lactate was sampled at multiple time points, actual individual peak values may occur outside this window [22]. This methodological limitation should be considered when interpreting [La]bmax.




Fig. (1). SmO₂ phases during incremental treadmill testing.

2.3. SRR

SRR was used to classify athletes based on their aerobic and anaerobic physiological attributes. It was calculated as the ratio of MSS to MAS, determined using the formula [10]:

$$SRR = MSS/MAS$$

MSS was obtained from the fastest 50-meter sprint trial recorded with laser timing gates. MAS was identified as the velocity at VO₂max (vVO₂max) during the treadmill test, based on breath-by-breath metabolic gas exchange analysis. The lowest treadmill speed at which VO₂max was maintained was designated as vVO₂max, following established methodology [6].

2.4. NIRS and SmO₂ Metrics

SmO₂ was continuously monitored using a Train.RED FYER NIRS device placed over the vastus lateralis. This non-invasive optical method calculates relative changes in total hemoglobin (tHb) by detecting the interplay between oxygenated and deoxygenated hemoglobin, using wavelengths at 760 nm (deoxyhemoglobin) and 850 nm (oxyhemoglobin). SmO₂ data were recorded at 10 Hz, consistent with the sampling rates reported in validation studies of wearable NIRS devices. Calibration was automatically done through the Train.RED app automatically during a one-minute initialization phase, as recommended by the manufacturer. Prior research has validated the reliability of the Train.RED FYER system for tracking hemoglobin saturation changes in sports science contexts [27]. To examine the muscle's oxygenation response during the incremental treadmill test, SmO₂ data were divided into four distinct phases (Fig. 1):

(1) **Baseline** - A five-minute seated rest period before the warm-up, establishing the individual's resting SmO₂ [25].

(2) **Test** - The entire duration of the treadmill protocol from onset to volitional exhaustion.

(3) **End-of-Test** - The final 30 seconds before the subject terminated the test, representing peak physiological strain [28].

(4) **Recovery** - The first minute immediately following exercise cessation, capturing initial reoxygenation patterns.

From these phases, four key SmO₂-derived metrics were calculated. ΔSmO₂ quantifies the magnitude of deoxygenation after an exercise bout by expressing the percent difference between baseline and End-of-Test SmO₂ values; calculated as the percent between these two values [16, 29]:
$$\Delta SmO_2 = \frac{Baseline\ SmO_2 - End\ of\ Test\ SmO_2}{Baseline\ SmO_2} \times 100\%.$$
 MdO₂ slope reflects the rate of SmO₂ decline at exercise onset and was obtained by fitting a segmented regression to the Test phase and extracting the linear slope up to the second major inflection point in the SmO₂-time curve, isolating the primary deoxygenation phase before plateau (Fig. 2). Several studies have shown that muscle deoxygenation often exhibits more than one breakpoint, reflecting different stages of oxygen extraction and metabolic demand during incremental exercise. By focusing on the interval between the start of exercise and this second inflection, we capture the linear portion of deoxygenation that best represents the upper limit of oxygen extraction before a subsequent plateau or slower rate of change [17, 30, 31]. For the IP metric, we chose

the first major breakpoint in the SmO_2 curve, identified via the same segmented regression analysis used for the MdO_2 slope (Fig. 3). The inflection marks the transition from predominantly aerobic to increasing anaerobic metabolism and corresponds closely with systemic lactate

thresholds (VT1/LT1), indicating when oxygen supply begins to lag behind demand [18]. Finally, the ROHS is defined as the linear rate of SmO_2 increase during the Recovery phase, with steeper slopes reflecting faster muscle reoxygenation driven by local perfusion and mitochondrial function (Fig. 4) [19].




Fig. (2). MdO_2 slope segmented regression analysis.




Fig. (3). Determination of IP using segmented regression analysis.




Fig. (4). Determination of ROHS post-maximal exercise.

2.5. Data Analysis

Statistical analyses were carried out in SPSS 26 (IBM, Armonk, USA) and R Studio 4.2.1 (R Core Team, Vienna, Austria). Variables were described using means \pm SD, medians, ranges, and kurtosis, and distribution normality was assessed using the Shapiro-Wilk test. To explore relationships among SmO₂ metrics, [La]bmax, and SRR, Pearson correlation coefficients were calculated and 95% confidence intervals were reported for each estimate. The alpha level was set a priori at <0.05 . Data trends and the strength of linear associations were visualized with scatterplots and segmented-regression charts. SmO₂ kinetics were examined by identifying key inflection points and computing deoxygenation and reoxygenation slopes. All procedures adhered to standard practices in sports science research. All figures were generated using Python (Matplotlib library).

3. RESULTS

3.1. Descriptive Statistics

A total of 11 NCAA Division I middle-distance runners

participated in this study. The average age of participants was 20.36 ± 1.57 years, with an average height of 178.12 ± 7.08 cm, and an average weight of 65.51 ± 6.62 kg. The body fat percentage averaged $7.56 \pm 1.78\%$, and the average weekly mileage reported was 48.18 ± 18.34 miles. Table 1 presents the descriptive statistics of key performance and physiological variables (MSS, MAS, SRR, VO2max, [La]bmax, Δ SmO₂, MdO₂, ROHS, and IP). The mean (\pm SD) VO2max was 68.09 ± 4.90 ml/kg/min, and the average SRR was 1.48 ± 0.16 , with values ranging from 1.34 to 1.73.

3.2. Correlation Heat Map

(Fig. 5) presents a Pearson correlation heat map summarizing the relationships among key physiological and performance metrics (MSS, MAS, SRR, VO2max, [La]bmax, Δ SmO₂, MdO₂, ROHS, and IP). Notably, SRR displayed a moderate positive correlation with Δ SmO₂ ($r = 0.66$, $p = 0.028$) and a moderate negative correlation with ROHS ($r = -0.64$, $p = 0.034$). In contrast, [La]bmax did not show a statistically significant relationship with SRR ($r = 0.25$, $p = 0.457$).


Table 1. Descriptive statistics of performance metrics.

	N	Minimum	Maximum	Mean	Std. Error	Std. Deviation	Kurtosis
MSS	11	28.20	35.30	30.60	0.62	2.04	1.68
MAS	11	18.10	21.85	20.59	0.36	1.19	0.46
SRR	11	1.34	1.73	1.48	0.05	0.16	-0.93
VO2max	11	60.23	75.60	68.09	1.48	4.90	-0.87
[La]bmax	11	6.10	19.60	11.76	1.35	4.46	-1.01

	N	Minimum	Maximum	Mean	Std. Error	Std. Deviation	Kurtosis
ΔSmO_2	11	27.23	52.89	34.19	2.15	7.13	4.83
MdO_2	11	-11.84	3.10	-4.59	1.29	4.27	-0.14
ROHS	11	43.20	78.45	62.99	3.44	11.42	-0.88
IP	11	6.12	8.67	7.66	0.23	0.77	0.05

Note: N = 11. MSS, maximal sprint speed (m/s); MAS, maximal aerobic speed (km/h); SRR, speed reserve ratio (MSS divided by MAS); $VO_2\text{max}$, maximal oxygen uptake ($\text{ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$); $[\text{La}]b\text{max}$, peak blood lactate concentration (mmol/L); ΔSmO_2 , percent change in muscle oxygen saturation from baseline to end of test; MdO_2 , muscle deoxygenation slope; ROHS, rate of muscle reoxygenation following exercise; IP, inflection point of SmO_2 curve during incremental exercise. Notably, $[\text{La}]b\text{max}$ ranged from 6.1 to 19.6 $\text{mmol}\cdot\text{L}^{-1}$, whereas SRR ranged only from 1.34 to 1.73, which may have reduced the power of correlations.

MSS

**Fig. (5).** Pearson correlation heat map.

Note: Warmer colors represent positive correlations and cooler colors represent negative correlations. Asterisks indicate statistical significance (* $p < 0.05$, ** $p < 0.01$).

3.3. Relationships Between SRR and Physiological Markers

As illustrated in Fig. (6A), ΔSmO_2 demonstrated a significant positive correlation with SRR ($r = 0.66$, $p = 0.028$). Athletes with higher SRR (*i.e.*, relatively more anaerobic-inclined) tended to show a larger percent change in muscle deoxygenation, whereas lower-SRR (more aerobic) runners exhibited smaller ΔSmO_2 values. Fig. (6B) shows the inverse relationship between ROHS

and SRR ($r = -0.64$, $p = 0.034$). Participants with lower SRR (suggesting stronger aerobic capacity) displayed faster reoxygenation rates, whereas higher-SRR individuals recovered more slowly post-exercise. Fig. (6C) shows the correlation between SRR, with the data points widely scattered, indicating substantial inter-individual variability. Although $[\text{La}]b\text{max}$ spanned a wide range across participants (6.1-19.6 mmol/L), no significant correlation emerged with SRR ($r = 0.25$, $p = 0.457$).




Fig. (6). (A-C) Correlation scatter plots.

Note: Scatter plots showing the relationships between speed reserve ratio (SRR) and key physiological variables. Panel A displays the correlation between SRR and percent change in muscle oxygen saturation (ΔSmO_2); Panel B shows SRR versus the reoxygenation slope (ROHS); and Panel C presents SRR versus peak blood lactate concentration ($[\text{La}]_{\text{bmax}}$). Each point represents an individual participant (N = 11), and dashed lines indicate the line of best fit.

4. DISCUSSION

This study aimed to determine whether SmO_2 -derived metrics and/or $[\text{La}]_{\text{bmax}}$ could serve as practical, noninvasive alternatives to the speed reserve ratio (SRR) for profiling the aerobic-anaerobic balance in NCAA Division I middle-distance runners. Of the SmO_2 metrics examined, only ΔSmO_2 and ROHS significantly correlated with SRR. Contrary to our hypothesis, ΔSmO_2 correlated positively with SRR ($r = 0.656$, $p = 0.028$), while ROHS supported our hypothesis with a negative correlation ($r = -0.644$, $p = 0.033$). These findings reveal a complex relationship between local oxygen utilization patterns and physiological profiles. IP, MdO_2 slope, and $[\text{La}]_{\text{bmax}}$ did not show significant correlations with SRR or with each other.

One of the key findings was the unanticipated positive correlation between ΔSmO_2 and SRR. A higher SRR indicates greater anaerobic capacity, characterized by a high maximal sprint speed relative to aerobic capacity. In contrast, a lower SRR reflects stronger aerobic capacity and endurance-oriented physiology. We expected that runners with lower SRR would exhibit larger ΔSmO_2 due to their skeletal muscles' enhanced ability to extract and utilize oxygen [32-34]. Instead, runners with higher SRR showed the greatest muscle deoxygenation during the maximal exercise. This pattern mirrors earlier research on sprint athletes performing 30-second Wingate tests under simulated hypoxia [16]. A plausible explanation for this observation involves differences in muscle fiber

composition and capillary structure between anaerobic and aerobic athletes. Athletes with higher anaerobic capacity (higher SRR) typically have more fast-twitch (Type II) fibers, characterized by lower capillary density and mitochondrial content compared to slow-twitch (Type I) fibers. During intense exercise, Type II fibers rapidly utilize oxygen, potentially exceeding the rate at which oxygen can be delivered to the sarcolemma through local microvasculature. This mismatch leads to pronounced muscle deoxygenation at the onset of exercise and may persist under conditions of maximal exertion due to ongoing oxygen supply-demand imbalance [27, 35]. By contrast, endurance training stimulates angiogenesis and enhances microvascular function in the skeletal muscle, supporting sustained oxygen delivery and blunting the SmO_2 drop during exercise [36-39]. Together, these physiological differences suggest that ΔSmO_2 reflects underlying muscle fiber composition and vascular characteristics, which may help explain its positive association with SRR by distinguishing athletes with more anaerobic versus aerobic physiological profiles.

We observed a strong negative correlation between ROHS and SRR ($r = -0.644$, $p = 0.033$), supporting our initial hypothesis that athletes with lower SRR recover muscle oxygenation more rapidly following a maximal exercise bout. Such rapid SmO_2 recovery likely reflects multiple endurance-related adaptations. As stated before, endurance training increases both capillary and mitochondrial density in skeletal muscle, which has been shown to significantly improve oxygen delivery and utilization during exercise and recovery periods [38, 39]. In addition, aerobic athletes tend to have increased myoglobin content, which can enhance intracellular oxygen diffusion, facilitating a faster return of SmO_2 level after exercise [40]. These adaptations could underpin the steeper ROHS slopes seen in the athletes with lower.

This study did not find any significant correlation with $[\text{La}]_{\text{bmax}}$ SRR or any SmO_2 metric, indicating that this metric alone cannot distinguish aerobic-anaerobic profiles in middle-distance runners. Although higher lactate levels reflect increased glycolytic activity during exercise, their absolute value and timing of reaching peak levels are influenced by individual factors, including buffering capacity, blood volume shifts, and training status [13]. Additionally, variability in lactate clearance rates can cause true peak lactate to occur outside the standard 3-to-7-minute postexercise sampling window used in this study [41, 42]. It is also possible that, within a homogenous cohort of well-trained middle-distance athletes, physiological profiles are so similar that $[\text{La}]_{\text{bmax}}$ lacks the sensitivity to detect subtle interindividual differences. While lactate testing remains a useful tool for understanding metabolic stress during exercise, its ability to fully profile middle-distance runners may be limited.

In addition, IP and MdO_2 slope did not significantly correlate with the SRR or the other metrics. Threshold metrics from NIRS measurements can vary considerably

due to factors such as sensor placement, treadmill protocol, and individual variability in metabolic transitions, limiting their reliability [43, 44]. Minor variations in the NIRS optode placement or changes in treadmill grade can substantially alter detected thresholds, highlighting the challenges of using these metrics to accurately profile aerobic-anaerobic balance in athletes.

Overall, these findings indicate that ΔSmO_2 and ROHS seem to be valid alternatives to the SRR and provide valuable insights into an athlete's physiology. These metrics may be incorporated into an athletic training program in order to understand an athlete's strengths and weaknesses, how they respond to training, and the race strategy that might suit them best. Systematic tracking of these metrics across training cycles may help coaches make adjustments in an athlete's exercise choice, intensity, and rest intervals in order to improve race performance while reducing the risk of overtraining.

Despite these promising insights, several limitations of our study must be acknowledged. First, the use of two different treadmill models (Cybex 751T and Woodway ELG) introduced potential variability in belt calibration, cushioning, and running surface characteristics, possibly influencing results despite standardization efforts. Second, our relatively small sample size ($n=11$) comprised only males, which restricts the generalizability of these findings to broader populations of middle-distance runners. Future research should include larger and more diverse cohorts, incorporating both male and female athletes across varying competitive levels, to enhance statistical power and clarify whether SmO_2 and $[\text{La}]_{\text{bmax}}$ responses differ significantly between subgroups. Third, high variability was found in multiple of the metrics, such as the MdO_2 and IP metrics. This raises concerns about reliability that should be considered when interpreting these measures. Fourth, fixed lactate sampling intervals (1-, 3-, 5-, and 7-minute post-exercise) might not have captured the true peak lactate concentration in all athletes due to individual differences in lactate response and clearance rates. Future studies should consider either continuous lactate monitoring or increasing the sampling intervals. Fifth, the absence of direct on-track performance data limits the practical application of the metrics. Future research should integrate these metrics with actual competition performance data to improve the predictive validity. Finally, the cross-sectional design of this study provides only a snapshot of athlete physiology and does not assess how interventions targeting ΔSmO_2 or ROHS affect performance over time. Prospective, intervention-based studies that manipulate training variables based on SmO_2 feedback could yield valuable insights into optimal training strategies tailored to individual aerobic and anaerobic profiles.

CONCLUSION

This study investigated whether SRR, SmO_2 metrics, and $[\text{La}]_{\text{bmax}}$ could collectively inform the aerobic-anaerobic profiles of NCAA Division I middle-

distance runners. Our findings indicate that SRR is a valuable composite measure reflecting an athlete's balance between sprinting and endurance capacities, with ΔSmO_2 positively correlating and ROHS negatively correlating with SRR. These correlations highlight the utility of portable NIRS technology for distinguishing athletes who rely heavily on anaerobic power (higher SRR, greater ΔSmO_2) from those who are better adapted aerobically (lower SRR, steeper ROHS). Conversely, $[\text{La}]_{\text{bmax}}$ alone did not significantly correlate with SRR or other physiological metrics.

Practically, coaches can leverage SmO_2 metrics during training to individualize workout intensities, optimize recovery intervals, and monitor physiological responses. Regular tracking of ΔSmO_2 and ROHS can assist in identifying appropriate training interventions, detecting early signs of fatigue, and preventing overtraining. Given the limitations associated with the small sample size and the absence of direct competition performance data, future research should seek to validate these findings in larger, more diverse athlete cohorts and explore longitudinal relationships between NIRS-derived metrics and actual race outcomes. Overall, this study supports integrating noninvasive SmO_2 monitoring into training programs to enhance athlete-specific training strategies and performance outcomes in middle-distance runners.

AUTHORS' CONTRIBUTIONS

The authors confirm their contribution to the paper as follows: J.A.C.: Study conception and design; B.K., A.K.: Data collection; E.E.S., K.A.E.: Analysis and interpretation of results; K.A.V.: Writing - reviewing and editing. All authors reviewed the results and approved the final version of the manuscript.

LIST OF ABBREVIATIONS

SRR	= Speed Reserve Ratio
SmO_2	= Skeletal Muscle Oxygenation
$[\text{La}]_{\text{bmax}}$	= Blood Lactate Concentration
ΔSmO_2	= Change in Skeletal Muscle Oxygenation
ROHS	= Reoxygenation Hemoglobin Slope
MSS	= Maximum Sprint Speed
MAS	= Maximum Aerobic Speed
NIRS	= Near-infrared Spectroscopy
MdO_2	= Muscle Deoxygenation
IP	= Inflection Point
tHb	= Total Hemoglobin

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

The project was approved by the California State University, Long Beach Institutional Review Board, for project number 1887311-4.

HUMAN AND ANIMAL RIGHTS

All procedures performed in studies involving human participants were in accordance with the ethical standards of institutional and/or research committee and with the 1975 Declaration of Helsinki, as revised in 2013.

CONSENT FOR PUBLICATION

Informed consent was obtained from all subjects who participated in the research.

AVAILABILITY OF DATA AND MATERIALS

The data and supportive information are available within the article.

STANDARDS OF REPORTING

STROBE guidelines were followed.

FUNDING

None.

CONFLICT OF INTEREST

Valenzuela Kevin A., is the Editorial Advisory Board member of The Open Sports Sciences Journal.

ACKNOWLEDGEMENTS

The authors would like to extend their sincere gratitude to the California State University, Long Beach Track and Field team for their invaluable support and participation during data collection. Their cooperation and dedication were instrumental in making this study possible.

DISCLOSURE

Part of this article has been previously published in the International Journal of Exercise Science Conference Proceedings as an abstract submitted to the Southwest American College of Sports Medicine Annual Conference (*International Journal of Exercise Science Conference Proceedings*: Vol. 14: Iss. 3, Article 92, November 2023).

REFERENCES

- [1] Spencer MR, Gastin PB. Energy system contribution during 200- to 1500-m running in highly trained athletes. *Med Sci Sports Exerc* 2001; 33(1): 157-62. <http://dx.doi.org/10.1097/00005768-200101000-00024> PMID: 11194103
- [2] Duffield R, Dawson B, Goodman C. Energy system contribution to 400-metre and 800-metre track running. *J Sports Sci* 2005; 23(3): 299-307. <http://dx.doi.org/10.1080/02640410410001730043> PMID: 15966348
- [3] Sandford GN, Stellingwerff T. "Question Your Categories": The misunderstood complexity of middle-distance running profiles with implications for research methods and application. *Front Sports Act Living* 2019; 1: 28. <http://dx.doi.org/10.3389/fspor.2019.00028> PMID: 33344952
- [4] Tanji F, Tsuji T, Shimazu W, Nabekura Y. Relationship between 800-m Running Performance and Aerobic and Anaerobic Energy Metabolism Capacities in Well-Trained Middle-Distance Runners. *Int J Sport Health Sci* 2018; 16(0): 70-6. <http://dx.doi.org/10.5432/ijshs.201724>
- [5] Thron M, Dükung P, Ruf L, Härtel S, Woll A, Altmann S. Assessing

anaerobic speed reserve: A systematic review on the validity and reliability of methods to determine maximal aerobic speed and maximal sprinting speed in running-based sports. *PLoS One* 2024; 19(1): 0296866. <http://dx.doi.org/10.1371/journal.pone.0296866> PMID: 38252665

[6] Billat LV. Interval training for performance: A scientific and empirical practice. Special recommendations for middle- and long-distance running. Part I: Aerobic interval training. *Sports Med* 2001; 31(1): 13-31. <http://dx.doi.org/10.2165/00007256-200131010-00002> PMID: 11219499

[7] Daniels J. 800 Meters. Daniels' Running Formula. Champaign, IL: Human Kinetics 2022; pp. 163-76. <http://dx.doi.org/10.5040/9781718219113.ch-011>

[8] Ingham SA, Whyte GP, Pedlar C, Bailey DM, Dunman N, Nevill AM. Determinants of 800-m and 1500-m running performance using allometric models. *Med Sci Sports Exerc* 2008; 40(2): 345-50. <http://dx.doi.org/10.1249/mss.0b013e31815a83dc> PMID: 18202566

[9] Wackerhage H, Schoenfeld BJ. Personalized, evidence-informed training plans and exercise prescriptions for performance, fitness and health. *Sports Med* 2021; 51(9): 1805-13. <http://dx.doi.org/10.1007/s40279-021-01495-w> PMID: 34143410

[10] Sandford GN, Laursen PB, Buchheit M. Anaerobic speed/power reserve and sport performance: Scientific basis, current applications and future directions. *Sports Med* 2021; 51(10): 2017-28. <http://dx.doi.org/10.1007/s40279-021-01523-9> PMID: 34398445

[11] Sandford GN, Kilding AE, Ross A, Laursen PB. Maximal sprint speed and the anaerobic speed reserve domain: The untapped tools that differentiate the world's best male 800 m runners. *Sports Med* 2019; 49(6): 843-52. <http://dx.doi.org/10.1007/s40279-018-1010-5> PMID: 30374943

[12] Ferri A, Adamo S, La Torre A, Marzorati M, Bishop DJ, Misericocchi G. Determinants of performance in 1,500-m runners. *Eur J Appl Physiol* 2012; 112(8): 3033-43. <http://dx.doi.org/10.1007/s00421-011-2251-2> PMID: 22179856

[13] Beneke R, Leithäuser RM, Ochentel O. Blood lactate diagnostics in exercise testing and training. *Int J Sports Physiol Perform* 2011; 6(1): 8-24. <http://dx.doi.org/10.1123/ijssp.6.1.8> PMID: 21487146

[14] Gupta S, Stanula A, Goswami A. Peak blood lactate concentration and its arrival time following different track running events in under-20 male track athletes. *Int J Sports Physiol Perform* 2021; 16(11): 1625-33. <http://dx.doi.org/10.1123/ijssp.2020-0685> PMID: 33848978

[15] Lacour JR, Bouvat E, Barthélémy JC. Post-competition blood lactate concentrations as indicators of anaerobic energy expenditure during 400-m and 800-m races. *Eur J Appl Physiol Occup Physiol* 1990; 61(3-4): 172-6. <http://dx.doi.org/10.1007/BF00357594> PMID: 2282899

[16] Oguri K, Fujimoto H, Sugimori H, Miyamoto K, Tachi T, Nagasaki S. Pronounced muscle deoxygenation during supramaximal exercise under simulated hypoxia in sprint athletes. *J Sports Sci Med* 2008; 7(4): 512-9. <http://dx.doi.org/10.5208/jssm.2008.512>

[17] Gasser B, Flück M, Frey WO, Valdivieso P, Spörri J. Association of gene variants for mechanical and metabolic muscle quality with cardiorespiratory and muscular variables related to performance in skiing athletes. *Genes* 2022; 13(10): 1798. <http://dx.doi.org/10.3390/genes13101798> PMID: 36292682

[18] Sendra-Pérez C, Encarnacion-Martinez A, Salvador-Palmer R, Murias JM, Priego-Quesada JI. Profiles of muscle-specific oxygenation responses and thresholds during graded cycling incremental test. *Eur J Appl Physiol* 2025; 125(1): 237-45. <http://dx.doi.org/10.1007/s00421-024-05593-1> PMID: 39259396

[19] Buchheit M, Ufland P. Effect of endurance training on performance and muscle reoxygenation rate during repeated-sprint running. *Eur J Appl Physiol* 2011; 111(2): 293-301. <http://dx.doi.org/10.1007/s00421-010-1654-9> PMID: 20872150

[20] Martin JS, Friedenreich ZD, Borges AR, Roberts MD. Acute effects of peristaltic pneumatic compression on repeated anaerobic exercise performance and blood lactate clearance. *J Strength Cond Res* 2015; 29(10): 2900-6. <http://dx.doi.org/10.1519/JSC.00000000000000928> PMID: 25756325

[21] Ghosh AK. Anaerobic threshold: Its concept and role in endurance sport. *Malays J Med Sci* 2004; 11(1): 24-36. PMID: 22977357

[22] Mavroudi M, Kabasaki A, Petridou A, Mougios V. Blood lactate and maximal lactate accumulation rate at three sprint swimming distances in highly trained and elite swimmers. *Sports* 2023; 11(4): 87. <http://dx.doi.org/10.3390/sports11040087> PMID: 37104161

[23] Bonaventura JM, Sharpe K, Knight E, Fuller KL, Tanner RK, Gore CJ. Reliability and accuracy of six hand-held blood lactate analysers. *J Sports Sci Med* 2015; 14(1): 203-14. PMID: 25729309

[24] ACSM's Guidelines for Exercise Testing and Prescription. (12th ED.), ACSM 2020.

[25] Haugen T, Seiler S, Sandbakk Ø, Tønnessen E. The training and development of elite sprint performance: An integration of scientific and best practice literature. *Sports Med Open* 2019; 5(1): 44. <http://dx.doi.org/10.1186/s40798-019-0221-0> PMID: 31754845

[26] Feldmann A, Schmitz R, Erlacher D. Near-infrared spectroscopy-derived muscle oxygen saturation on a 0% to 100% scale: Reliability and validity of the Moxy Monitor. *J Biomed Opt* 2019; 24(11): 1-11. <http://dx.doi.org/10.1117/1.JBO.24.11.115001> PMID: 31741352

[27] Goodwin ML, Harris JE, Hernández A, Gladden LB. Blood lactate measurements and analysis during exercise: A guide for clinicians. *J Diabetes Sci Technol* 2007; 1(4): 558-69. <http://dx.doi.org/10.1177/193229680700100414> PMID: 19885119

[28] Victorio M, Dieffenderfer J, Songkakul T, Willeke J, Bozkurt A, Pozdin VA. Wearable wireless functional near-infrared spectroscopy system for cognitive activity monitoring. *Biosensors* 2025; 15(2): 92. <http://dx.doi.org/10.3390/bios15020092> PMID: 39996994

[29] Perrey S, Quaresima V, Ferrari M. Muscle oximetry in sports science: An updated systematic review. *Sports Med* 2024; 54(4): 975-96. <http://dx.doi.org/10.1007/s40279-023-01987-x> PMID: 38345731

[30] Paquette M, Bieuzen F, Billaut F. The effect of HIIT vs. SIT on muscle oxygenation in trained sprint kayakers. *Eur J Appl Physiol* 2021; 121(10): 2743-59. <http://dx.doi.org/10.1007/s00421-021-04743-z> PMID: 34145486

[31] Yoge A, Arnold J, Nelson H, et al. The effect of severe intensity bouts on muscle oxygen saturation responses in trained cyclists. *Front Sports Act Living* 2023; 5: 1086227. <http://dx.doi.org/10.3389/fspor.2023.1086227> PMID: 36909360

[32] Boone J, Koppo K, Barstow TJ, Bouckaert J. Pattern of deoxy[Hb + Mb] during ramp cycle exercise: Influence of aerobic fitness status. *Eur J Appl Physiol* 2009; 105(6): 851-9. <http://dx.doi.org/10.1007/s00421-008-0969-2> PMID: 19130074

[33] Yoge A, Arnold J, Clarke D, Guenette JA, Sporer BC, Koehle MS. Comparing the respiratory compensation point with muscle oxygen saturation in locomotor and non-locomotor muscles using wearable nirs spectroscopy during whole-body exercise. *Front Physiol* 2022; 13: 818733. <http://dx.doi.org/10.3389/fphys.2022.818733> PMID: 35431982

[34] Ross M, Kargl CK, Ferguson R, Gavin TP, Hellsten Y. Exercise-induced skeletal muscle angiogenesis: Impact of age, sex, angiocrine and cellular mediators. *Eur J Appl Physiol* 2023; 123(7): 1415-32. <http://dx.doi.org/10.1007/s00421-022-05128-6> PMID: 36715739

[35] Prefaut C, Durand F, Mucci P, Caillaud C. Exercise-induced arterial hypoxaemia in athletes: A review. *Sports Med* 2000; 30(1): 47-61.

<http://dx.doi.org/10.2165/00007256-200030010-00005> PMID: 10907757

[36] Dong G, Wang Y. Effect of physiological exercise on increasing the maximum oxygen uptake of skeletal muscle. *Rev Bras Med Esporte* 2021; 27(7): 710-3.
http://dx.doi.org/10.1590/1517-8692202127072021_0352

[37] Pringle JSM, Doust JH, Carter H, et al. Oxygen uptake kinetics during moderate, heavy and severe intensity 'submaximal' exercise in humans: The influence of muscle fibre type and capillarisation. *Eur J Appl Physiol* 2003; 89(3): 289-300.
<http://dx.doi.org/10.1007/s00421-003-0799-1> PMID: 12736837

[38] Abernethy PJ, Thayer R, Taylor AW. Acute and chronic responses of skeletal muscle to endurance and sprint exercise. A review. *Sports Med* 1990; 10(6): 365-89.
<http://dx.doi.org/10.2165/00007256-199010060-00004> PMID: 2291032

[39] Merlet AN, Messonnier LA, Coudy-Gandilhon C, et al. Beneficial effects of endurance exercise training on skeletal muscle microvasculature in sickle cell disease patients. *Blood* 2019; 134(25): 2233-41.
<http://dx.doi.org/10.1182/blood.2019001055> PMID: 31742587

[40] Andersen P, Henriksson J. Capillary supply of the quadriceps femoris muscle of man: Adaptive response to exercise. *J Physiol* 1977; 270(3): 677-90.
<http://dx.doi.org/10.1113/jphysiol.1977.sp011975> PMID: 198532

[41] Vernillo G, Brighenti A, Limonta E, et al. Effects of ultratrail running on skeletal-muscle oxygenation dynamics. *Int J Sports Physiol Perform* 2017; 12(4): 496-504.
<http://dx.doi.org/10.1123/ijssp.2015-0745> PMID: 27617750

[42] Richardson RS, Noyszewski EA, Kendrick KF, Leigh JS, Wagner PD. Myoglobin O₂ desaturation during exercise. Evidence of limited O₂ transport. *J Clin Invest* 1995; 96(4): 1916-26.
<http://dx.doi.org/10.1172/JCI118237> PMID: 7560083

[43] Foster C, Lucia A. Running economy. *Sports Med* 2007; 37(4): 316-9.
<http://dx.doi.org/10.2165/00007256-200737040-00011> PMID: 17465597

[44] Sendra-Pérez C, Sanchez-Jimenez JL, Marzano-Felisatti JM, Encarnación-Martínez A, Salvador-Palmer R, Priego-Quesada JI. Reliability of threshold determination using portable muscle oxygenation monitors during exercise testing: A systematic review and meta-analysis. *Sci Rep* 2023; 13(1): 12649.
<http://dx.doi.org/10.1038/s41598-023-39651-z> PMID: 37542055

DISCLAIMER: The above article has been published, as is, ahead-of-print, to provide early visibility but is not the final version. Major publication processes like copyediting, proofing, typesetting and further review are still to be done and may lead to changes in the final published version, if it is eventually published. All legal disclaimers that apply to the final published article also apply to this ahead-of-print version.