

The Addition of a Video Game to Stationary Cycling: The Impact on Energy Expenditure in Overweight Children

Bryan L. Haddock*, Shannon R. Siegel and Linda D. Wikin

Department of Kinesiology, California State University, San Bernardino

Abstract: *Introduction:* The prevalence of obesity in children has reached epidemic proportions with over 37% of children aged 6-11 years in the U.S. being classified as “at risk for overweight” or “overweight.” Utilization of active video games has been proposed as one possible mechanism to help shift the tide of the obesity epidemic.

Purpose: The purpose of this study was to determine if riding a stationary bike that controlled a video game would lead to significantly greater energy expenditure than riding the same bike without the video game connected.

Methods: Twenty children, 7-14 years old, with a BMI classification of “at risk for overweight” or “overweight” participated in this study. Following familiarization, energy expenditure was evaluated while riding a stationary bike for 20 minutes. One test was performed without the addition of a video game and one test with the bike controlling the speed of a car on the video game.

Results: Oxygen consumption and energy expenditure were significantly elevated above baseline in both conditions. Energy expenditure was significantly higher while riding the bike as it controlled the video game (4.4 ± 1.2 Kcal·min⁻¹) than when riding the bike by itself (3.7 ± 1.1 Kcal·min⁻¹) ($p < 0.05$). Perceived exertion was not significantly different between the two sessions ($p > 0.05$).

Conclusion: Using a stationary bike to control a video game led to greater energy expenditure than riding a stationary bike without the video game and without a related increase in perceived exertion.

Key Words: Obesity, Oxygen consumption, Kcal, BMI.

INTRODUCTION

The prevalence of obesity in children has continued to increase in recent years. According to the Center for Disease Control’s (CDC) classification system, the prevalence of children in the United States, aged 6-11 years, classified as “overweight” has reached 18.8%, while the prevalence of those classified as either “overweight” or “at risk for overweight” is at 37.2% [1, 2]. Although there is some evidence that the increased prevalence of obesity is leveling off in children, the prevalence is still considered excessively high [3]. This increased prevalence of overweight in children has become a global epidemic [4-9] and has been shown to be associated with multiple cardiovascular risk factors, including higher levels of cholesterol, triglycerides, systolic and diastolic blood pressure, and fasting insulin [10].

Children spend a large amount of time watching television and playing computer games [11]. The time children spend on these tasks has often been blamed for the increased prevalence of overweight in children [12-19]. Previous efforts to reduce the amount of screen time (television and video games) in children have proven to be successful in

helping children decrease their weight and improve their body composition [20, 21].

In recent years, interest has risen as to the efficacy of interactive video games (exergaming). Several studies have examined the level of energy expenditure while playing these exergames [22-26]. The level of energy expenditure has been shown to increase significantly above resting levels [22-24], or above the level of energy expenditure while playing sedentary video games [25, 26] and watching television [25]. Generally these studies have shown that energy expenditure will increase over resting levels to an intensity that is similar to moderate intensity exercise, with a VO_2 during activity of approximately 14-24 ml·kg⁻¹·min⁻¹.

A study of wheelchair bound athletes found that utilizing an arm ergometer to control a video game increased energy expenditure compared to utilizing the same arm ergometer without the video game interaction. This enhanced energy expenditure was found without a corresponding increase in perceived exertion [27]. No published research has compared the energy expenditure of overweight children while performing a traditional exercise (e.g., stationary cycling) to their energy expenditure while exercising on the same equipment with interactive gaming. Therefore, the purpose of this study was to determine if adding an interactive video game component to traditional cycle ergometry would impact the energy expenditure and rating of perceived exertion (RPE) in overweight children.

*Address correspondence to this author at the Department of Kinesiology, California State University, San Bernardino; Tel: (909)537-5359; Fax: (909)537-7085; E-mail: bhaddock@csusb.edu

METHODS

Subjects

Twenty children (13 boys, 7 girls), between the ages of 7 and 14 years participated in this study. The participants were recruited from a local weight management program and were generally sedentary. Each participant's height and weight were measured with a portable stadiometer (SECA 214; Hamburg, Germany) and calibrated digital scale (TANITA; Arlington Heights, Illinois), to the nearest 0.5 cm and 0.1 kg respectively. All participants had a Body Mass Index (BMI) that put them at the 85th percentile or above for their age and sex according to the CDC guidelines.[1] All participants were free of diabetes, thyroid disorders, heart disease, or hypertension. The research protocol was approved by the Institutional Review Board (IRB) of the California State University, San Bernardino. Child assent and written parental informed consent was obtained prior to participation.

Procedures

Each participant was scheduled for three appointments: (1) familiarization with the equipment, (2) testing session #1, and (3) testing session #2. At least one parent was required to be in attendance at each session.

At the familiarization session, each participant was shown the portable metabolic cart (Cosmed K4b² – Rome, Italy) including the mask and heart rate (HR) monitor that they would be wearing during the two testing sessions. Each participant was then allowed to ride on the cycle ergometer (CatEye™ Gamebike - USA) and play the games that would be utilized in the study. Participants were allowed to play the games during this session for as long as they wanted. All of the participants had previous experience with the ergometer, having utilized the system in their weight management program. The video game utilized for this study was Disney's "Cars" (Pixar Entertainment) on a PlayStation®2 console.

For each testing session, the children were instructed not to eat for a minimum of two hours prior to their appointment time, although water was allowed. All testing was done in a private room so that only the participant, parents, and researchers were present. Upon arriving, the participants were reminded of the testing procedures. The participants were then connected to the calibrated portable metabolic cart and HR monitor. The O₂ and CO₂ sensors were calibrated prior to each test with a known gas concentration of 16% O₂ and 5% CO₂. In addition, volume calibration was performed with a 3-Liter Syringe. Participants were then seated in a comfortable chair. Breath by breath oxygen consumption (VO₂) and carbon dioxide production (VCO₂) were collected for 10 minutes, with the last 5-minutes of collection averaged for determination of baseline energy expenditure.

After collection of the baseline data, participants stepped up to the cycle ergometer for one of two testing sessions: (1) a 20-minute exercise session with no video game (NV) or (2) a 20-minute exercise session with the cycle ergometer controlling the video game (V). Breath by breath VO₂ and CO₂ were measured continuously over the 20-minute session to determine energy expenditure. The order of the two testing sessions was randomized and counterbalanced. The testing days were separated by a minimum of two days so that the participants would not experience left over fatigue from the

prior test. A couple of participants were tested a full week later, with no significant change in their activity in between sessions. The ergometer seat height was adjusted to the proper height with the same seat height utilized for both exercise sessions. The resistance on the flywheel was standardized for both tests. Once seated on the ergometer, the participant was again reminded of the procedure. The participants were asked to ride the bike for 20 minutes, with the stated goal to get as much exercise in as possible, but they were free to slow down and rest at any time they desired. In order to keep a consistent pattern of encouragement, the researchers gave positive feedback every two minutes during both exercise sessions. The participants would be told they were doing well and asked how they were feeling. Heart rate, oxygen consumption, and carbon dioxide production were measured continuously throughout the test in order to determine energy expenditure. At the end of the 20 minutes of riding, the participants stepped off of the ergometer and walked around the room to cool down or sat in a chair and rocked their feet to prevent blood pooling. During the cool-down period, the participants were asked to rate their perceived level of exertion based on the OMNI perceived exertion scale (RPE) (See Fig. 1) that was specifically designed for children [28].

All data were analyzed using SPSS 14.0. Descriptive data (Mean ± SD) were calculated for the age in years, height in centimeters, weight in kilograms, BMI (kg/m²), and energy expenditure at baseline and during the exercise time. A paired sample *t*-test was utilized to determine differences between NV and V conditions. A one-way ANOVA was utilized to determine if any differences existed between males and females. Pearson product-moment correlations were performed on continuous variables to determine whether any significant relationships existed.

RESULTS

Descriptive data of the participants are provided in Table 1. All participants had a BMI that was above the 85th percentile for their age and sex based on the CDC guidelines.[1] No significant differences existed between boys and girls on baseline characteristics (*p*>0.05). In addition there were no significant differences at baseline for the two different exercise conditions, V and NV (*p*>0.05).

Oxygen consumption (ml•kg⁻¹•min⁻¹) was significantly elevated above baseline conditions in both exercise sessions (*p*<0.001). Table 2 summarizes the difference between the two exercise conditions. The net increase in energy expenditure (Gross energy Kcal•min⁻¹ minus resting Kcal•min⁻¹) was significantly higher (*p*<0.01) in the V session (4.4 ± 1.2 Kcal/min) than the NV session (3.7 ± 1.1 Kcal•min⁻¹). Participants expended a total of 98.7 ± 27.2 Kcal during the NV session and 113.2 ± 31.6 Kcal during the V session.

The average HR was not significantly different in the two testing sessions. The average HR during the NV session reached 142.4 ± 18.8 bpm, or 68% of age predicted max HR. During the V session, the average HR was 146.0 ± 21.4 bpm, or 70% of age predicted max HR. The peak HR achieved during the entire session was significantly higher in the V session than the NV session (*p*<0.01).

The average RPE was not significantly different between the two sessions (3.6 ± 2.3 in the NV session and 3.2 ± 2.8 in




Fig. (1). Validated OMNI Rating of Perceived Exertion Scale.

Table 1. Baseline Descriptive Data (Mean \pm SD)

| | Boys (n=13) | Girls (n=7) | Pooled Data (n=20) |
|--------------------------|------------------|-----------------|--------------------|
| Age (yrs.) | 10.8 \pm 2.3 | 11.2 \pm 2.3 | 10.9 \pm 2.2 |
| Height (cm.) | 151.2 \pm 17.3 | 153.9 \pm 7.4 | 152.1 \pm 14.5 |
| Weight (kg.) | 72.8 \pm 39.6 | 76.7 \pm 14.9 | 74.1 \pm 32.6 |
| BMI (kg/m ²) | 29.2 \pm 9.2 | 34.0 \pm 5.5 | 30.9 \pm 8.3 |

the V session; $p > 0.05$). RPE was significantly correlated to the participant's BMI following the NV session ($r = 0.55$; $p < 0.05$), but not following the V session ($r = 0.29$; $p > 0.05$). BMI was significantly correlated to the energy expenditure in both the V ($r = 0.78$) and NV ($r = 0.70$) exercise sessions ($p < 0.001$).

DISCUSSION

It is clear that the prevalence of children who are classified as "overweight" or "at risk for overweight" has reached a level that requires the examination of creative ways to combat this trend. Although the cause of the increased prevalence of overweight in children is multifaceted, a lack of adequate physical activity is accepted as one of these causes [29]. Increasing physical activity through exergaming has been postulated as a potential mechanism for increasing physical activity in children. Previous research has demonstrated that interactive video games can lead to a moderate increase in energy expenditure [22-26]. The level of energy expenditure in the current study (4.9 ± 1.4 Kcal \cdot min⁻¹ for the NV session and 5.7 ± 1.6 Kcal \cdot min⁻¹ for the V session) is similar to what might be expected with leisurely bike riding

[30]. Unlike the majority of the previously published work, the current study examined the use of exergaming in overweight children. Only one of the previous studies examined the response in overweight children and the children were a little older (13.5 ± 3.3 yrs) and had a lower mean BMI (27.4 ± 3.3 kg/m²) [23]. The overweight participants in this previous study expended an average of 4.6 ± 1.3 Kcal \cdot min⁻¹ while playing a dance simulation game for 12 minutes. The participants in the current study (10.9 ± 2.2 years, 30.9 ± 8.3 kg/m²) expended a greater amount of energy both with and without the utilization of the video game.

As seen in Table 2, the participants increased their oxygen consumption and energy expenditure above baseline in both exercise conditions. However, the addition of a video game to the stationary biking condition significantly increased energy expenditure ($p < 0.01$). When examining the total energy expended over the 20 minutes of exercise, the participants metabolized an extra 14.5 Kcal in the V session. Although statistically significant, this extra expended energy is probably of minimal clinical significance, unless the participants would be more likely to perform the activity with the video game and/or exercise for a longer period of time.

Table 2. Physiological Effect of Exercise: With and Without Video Games (Mean \pm SD; n=20)

| Variable | No Video Mean \pm SD | Video Mean \pm SD | t-value |
|---|------------------------|---------------------|---------------------|
| Resting Values | | | |
| H.R. (beats/min) | 90.9 \pm 13.4 | 91.5 \pm 14.5 | -0.157 |
| VO ₂ (ml•kg ⁻¹ •min ⁻¹) | 3.5 \pm 0.8 | 3.6 \pm 0.9 | -0.446 |
| RER | 0.92 \pm 0.14 | 0.94 \pm 0.29 | -0.369 |
| Kcal/min | 1.3 \pm 0.6 | 1.3 \pm 0.5 | -0.328 |
| Exercise Values | | | |
| Average HR (beats/min) | 142.4 \pm 18.8 | 146.0 \pm 21.4 | -0.889 |
| Max HR (beats/min) | 159.8 \pm 22.8 | 173.3 \pm 18.6 | -2.973 [†] |
| Average VO ₂ (ml•kg ⁻¹ •min ⁻¹) | 14.7 \pm 4.3 | 16.8 \pm 4.2 | -3.175 [†] |
| Peak VO ₂ (ml•kg ⁻¹ •min ⁻¹) | 19.3 \pm 5.7 | 21.9 \pm 6.2 | -2.815* |
| Average RER | 0.88 \pm 0.06 | 0.92 \pm 0.11 | -1.847 |
| Kcal/min | 4.9 \pm 1.4 | 5.7 \pm 1.6 | -3.240 [†] |
| Total Kcal (20 min.) | 98.7 \pm 27.2 | 113.2 \pm 31.6 | -3.238 [†] |
| RPE | 3.6 \pm 2.3 | 3.2 \pm 2.8 | 0.666 |

* $p < 0.05$; [†] $p < 0.01$.

A previous study has demonstrated that overweight children are likely to exercise longer if they have something to distract them from their exercise session [31]. Therefore, it would be expected that the addition of an interactive video game to a traditional exercise (stationary cycling in the current study) would distract overweight children from their perceived effort. The current study seems to bear this out. As shown in Table 2, the energy expenditure in the V session was significantly higher than in the NV session. However, the RPE was not significantly greater and in fact tended to be lower in the V session (3.6 \pm 2.3 in the NV session and 3.2 \pm 2.8 in the V session). It would appear likely that children would be more likely to exercise regularly and for a longer period of time while playing a game compared to riding a stationary bike without the interactive video game component. However, a study comparing the reinforcing value of interactive video games found that an interactive bike game was no more reinforcing than biking alone, whereas, interactive dance was more reinforcing [32]. A further evaluation of the utilization of interactive video games to increase the amount of time children will exercise is warranted.

The participants in this study exercised at an intensity that raised their HR to an average of 142.4 \pm 18.8 beats per minute (68% of age predicted max HR) and 146.0 \pm 21.4 beats per minute (70% of age predicted max HR) for the NV and V sessions respectively. This level of exertion is classified as moderate intensity exercise and therefore in line with what would be recommended to children [33]. Likewise, the oxygen consumption is 14.7 \pm 4.3 ml•kg⁻¹•min⁻¹ in the NV session and 16.8 \pm 4.2 ml•kg⁻¹•min⁻¹ in the V session. This is equal to 4.2 and 4.8 METS for the NV and V sessions respectively. This is similar to leisurely bike ride according to the compendium of physical activities [30].

CONCLUSION

Adding an active video game component to standard stationary biking led to a modest increase in energy expenditure

in this group of overweight children. However, this modest increase in energy expenditure occurred without a related increase in perceived exertion. Therefore, adding an active video game component to traditional exercise such as stationary cycling could lead to an increase in total energy expenditure, especially if the video game made the children more likely to exercise or to exercise for a longer period of time.

ACKNOWLEDGEMENTS

This study was supported in part by the National Institutes of Health (NIH grants # GM073842; HD052368; MD002722), and by the United States Department of Agriculture (USDA grant # 2008-55215-18837).

REFERENCES

- [1] CDC. Growth Charts: United States. 2000, May 30 [Accessed date July 1 2008] [cited; Available from: <http://www.cdc.gov/nchs/about/major/nhanes/growthcharts>]
- [2] Ogden CL, Carroll MD, Curtin LR, McDowell MA, Tabak CJ, Flegal KM. Prevalence of overweight and obesity in the United States, 1999-2004. *JAMA* 2006; 295: 1549-55.
- [3] Ogden CL, Carroll MD, Flegal KM. High body mass index for age among US children and adolescents, 2003-2006. *JAMA* 2008; 299: 2401-5.
- [4] Obesity: preventing and managing the global epidemic. Report of a WHO consultation. World Health Organization technical report series 2000; 894: 1-253.
- [5] Raymond SU, Leeder S, Greenberg HM. Obesity and cardiovascular disease in developing countries: a growing problem and an economic threat. *Curr Opin Clin Nutr Metab Care* 2006; 9: 111-6.
- [6] Reilly JJ, Dorosty AR. Epidemic of obesity in UK children. *Lancet* 1999; 354: 1874-5.
- [7] Rossner S. Obesity: the disease of the twenty-first century. *Int J Obes Relat Metab Disord* 2002; 26 (Suppl 4): S2-4.
- [8] Rudolf MC, Sahota P, Barth JH, Walker J. Increasing prevalence of obesity in primary school children: cohort study. *BMJ* 2001; 322: 1094-5.
- [9] Seidell JC, Verschuren WM, Kromhout D. Prevalence and trends of obesity in The Netherlands 1987-1991. *Int J Obes Relat Metab Disord* 1995; 19: 924-7.

- [10] Freedman DS, Dietz WH, Srinivasan SR, Berenson GS. The relation of overweight to cardiovascular risk factors among children and adolescents: the Bogalusa Heart Study. *Pediatrics* 1999; 103: 1175-82.
- [11] Marshall SJ, Gorely T, Biddle SJ. A descriptive epidemiology of screen-based media use in youth: a review and critique. *J Adolesc* 2006; 29: 333-49.
- [12] Andersen RE, Crespo CJ, Bartlett SJ, Cheskin LJ, Pratt M. Relationship of physical activity and television watching with body weight and level of fatness among children: results from the third national health and nutrition examination survey. *JAMA* 1998; 279: 938-42.
- [13] Crespo CJ, Smit E, Troiano RP, Bartlett SJ, Macera CA, Andersen RE. Television watching, energy intake, and obesity in US children: results from the third national health and nutrition examination survey, 1988-1994. *Arch Pediatr Adolesc Med* 2001; 155: 360-5.
- [14] Dietz WH, Jr, Gortmaker SL. Do we fatten our children at the television set? Obesity and television viewing in children and adolescents. *Pediatrics* 1985; 75: 807-12.
- [15] Gortmaker SL, Must A, Sobol AM, Peterson K, Colditz GA, Dietz WH. Television viewing as a cause of increasing obesity among children in the United States, 1986-1990. *Arch Pediatr Adolesc Med* 1996; 150: 356-62.
- [16] Hesketh K, Wake M, Graham M, Waters E. Stability of television viewing and electronic game/computer use in a prospective cohort study of Australian children: relationship with body mass index. *Int J Behav Nut Phys Act* 2007; 4: 60.
- [17] Janz KF, Levy SM, Burns TL, Torner JC, Willing MC, Warren JJ. Fatness, physical activity, and television viewing in children during the adiposity rebound period: the iowa bone development study. *Prev Med* 2002; 35: 563-71.
- [18] Stettler N, Signer TM, Suter PM. Electronic games and environmental factors associated with childhood obesity in Switzerland. *Obes Res* 2004; 12: 896-903.
- [19] Vandewater EA, Shim MS, Caplovitz AG. Linking obesity and activity level with children's television and video game use. *J Adolesc* 2004; 27: 71-85.
- [20] Robinson TN, Killen JD, Kraemer HC, *et al.* Dance and reducing television viewing to prevent weight gain in African-American girls: the Stanford GEMS pilot study. *Ethn Dis* 2003;13: S65-77.
- [21] Robinson TN. Reducing children's television viewing to prevent obesity: a randomized controlled trial. *JAMA* 1999; 282: 1561-7.
- [22] Tan B, Aziz AR, Chua K, Teh KC. Aerobic demands of the dance simulation game. *Int J Sports Med* 2002; 23:125-9.
- [23] Unnithan VB, Houser W, Fernhall B. Evaluation of the energy cost of playing a dance simulation video game in overweight and non-overweight children and adolescents. *Int J Sports Med* 2006; 27: 804-9.
- [24] Maddison R, Mhurchu CNI, Jull A, Jiang Y, Prapavessis H, Rodgers A. Energy expended playing video console games: an opportunity to increase children's physical activity? *Pediatr Exerc Sci* 2007; 19: 334-43.
- [25] Lanningham-Foster L, Jensen TB, Foster RC, *et al.* Energy expenditure of sedentary screen time compared with active screen time for children. *Pediatrics* 2006;118: e1831-5.
- [26] Graves L, Stratton G, Ridgers ND, Cable NT. Comparison of energy expenditure in adolescents when playing new generation and sedentary computer games cross sectional study. *BMJ* 2007; 335: 1282-4.
- [27] Fitzgerald SG, Cooper RA, Thorman T, Cooper R, Guo S, Boninger ML. The GAME(Cycle) exercise system: comparison with standard ergometry. *J Spinal Cord Med* 2004; 27: 453-9.
- [28] Utter AC, Robertson RJ, Nieman DC, Kang J. Children's OMNI Scale of Perceived Exertion: walking/running evaluation. *Med Sci Sports Exerc* 2002; 34: 139-44.
- [29] Anderson PM, Butcher KE. Childhood obesity: trends and potential causes. *Future Child* 2006; 16: 19-45.
- [30] Ainsworth BE, Haskell WL, Whitt MC, *et al.* Compendium of physical activities: an update of activity codes and MET intensities. *Med Sci Sports Exerc* 2000; 32: S498-504.
- [31] De Bourdeaudhuij I, Crombez G, Deforche B, Vinaimont F, Debode P, Bouckaert J. Effects of distraction on treadmill running time in severely obese children and adolescents. *Int J Obes Relat Metab Disord* 2002; 26: 1023-9.
- [32] Epstein LH, Beecher MD, Graf JL, Roemmich JN. Choice of interactive dance and bicycle games in overweight and nonoverweight youth. *Ann Behav Med* 2007; 33: 124-31.
- [33] Medicine ACoS. ACSM's guidelines for exercise testing and prescription. 7th ed. Philadelphia: Lippincott Williams & Wilkins 2006.

Received: October 20, 2008

Revised: February 4, 2009

Accepted: February 4, 2009

© Haddock *et al.*; Licensee Bentham Open.This is an open access article licensed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted, non-commercial use, distribution and reproduction in any medium, provided the work is properly cited.